

Classical and Quantum Solutions of 2+1 Dimensional Gravity under De Broglie-Bohm Interpretation

M. Kenmoku ¹, T. Matsuyama ², R. Sato ³ and S. Uchida ⁴

¹ *Department of Physics, Nara Women's University, Nara 630-8506, Japan*

² *Department of Physics, Nara University of Education, Takabatake-cho, Nara 630-8528, Japan*

^{3 4} *Graduate School of Human Culture, Nara Women's University, Nara 630-8506, Japan*

Abstract

We have studied classical and quantum solutions of 2+1 dimensional Einstein gravity theory. Quantum theory is defined through the local conserved angular momentum and mass operators in the case of spherically symmetric space-time. The de Broglie-Bohm interpretation is applied for the wave function and we derive the differential equations for metrics. Metrics including quantum effect are obtained in solving these equations and we compare them with classical metrics. Especially the quantum effect on the closed de Sitter universe is evaluated quantitatively.

PACS numbers: 98.80.Hw, 03.65.Bz, 04.60.Kz, 04.60.-m

¹ kenmoku@phys.nara-wu.ac.jp

² matsuyat@nara-edu.ac.jp

³ reika@phys.nara-wu.ac.jp

⁴ satoko@phys.nara-wu.ac.jp

1 Introduction

Recently 2+1 dimensional gravity theory, especially AdS_3 has been studied extensively [1, 2, 3]. It was shown that the theory is equivalent to the 2+1 Chern-Simons theory [4] and has been investigated to understand the black hole thermodynamics, i.e. Hawking temperature [5] and others. The 2+1 dimensional AdS supergravity theories were investigated in the Chern-Simons form [6]. Topologically massive gravity, which is the sum of Einstein and Chern-Simons terms, was also analyzed in the canonical form [7]. The purpose of this paper is to investigate the canonical formalism of the original 2+1 dimensional Einstein gravity theory instead of the Chern-Simons theory. For the spherically symmetric space-time, local conserved quantities (mass and angular momentum) are introduced and using them canonical quantum theory is defined. Constraints are imposed on state vectors and solved analytically. In order to extract the physical meaning of the wave function, we adopt the de Broglie-Bohm interpretation [8, 9, 10] and derive the differential equations for the metrics, which include the quantum effect.

After fixing the gauge choice, special solutions of the metrics are obtained. Especially the quantum effect on the closed de Sitter universe is obtained quantitatively. It is interesting to note that the birth of universe appears as the real time tunneling instead of the imaginary time tunneling in the WKB approximation by Vilenkin [11] and in the path integral method by Hartle and Hawking [12]. The strategy to obtain the solution is followed by our previous work in the 3+1 dimensional gravity [13]. The new developments in this paper are to take into account a general form of metrics and to make a quantitative evaluation to

quantum effects in the 2+1 dimensional spherically symmetric gravity.

This paper is organized as follows. In section 2, we review the canonical formalism of the Einstein theory with a cosmological constant in spherically symmetric space-time. We introduce the local conservation quantities, the angular momentum J and the mass function M . In section 3, the canonical quantization is presented. Using some operator ordering the quantum theory could be constructed and the analytic solutions could be obtained. In section 4, we adopt the de Broglie-Bohm interpretation in order to extract geometrical information from the analytic solutions. Some examples of the comparison between classical limit and classical solutions which have already been known are also exhibited. In section 5, we try to evaluate the finite quantum effect for closed de Sitter universe. Summary and concluding remark are given in the final section.

2 Canonical Formalism

We start to consider the Einstein-Hilbert action with cosmological constant λ in 2+1 dimensional space-time,

$$I = \frac{1}{16\pi G_2} \int d^3x \sqrt{-^{(3)}g} (^{(3)}R - 2\lambda). \quad (1)$$

The gravitational constant in 2+1 dimensions is set to $G_2 = 1/4$ in the following. The metrics in polar coordinates are expressed in ADM decomposition [14] as

$$ds^2 = -N^2 dt^2 + \Lambda^2 (dr + N^r dt)^2 + R^2 (d\phi + N^\phi dt)^2 + 2C(dr + N^r dt)(d\phi + N^\phi dt), \quad (2)$$

where all metrics are assumed to be function of time t and radial coordinate r . In the following, dot and dash denotes the derivative with respect to t and r .

The action in the canonical formalism is in the form,

$$I = \int dt dr [P_\Lambda \dot{\Lambda} + P_R \dot{R} + P_C \dot{C} - (NH + N^r H_r + N^\phi H_\phi)] - \int dt dr \left([(\Lambda P_\Lambda + C P_C) N^r]' + \left[\left(\frac{C}{\Lambda} P_\Lambda + R^2 P_C \right) N^\phi \right]' \right), \quad (3)$$

where canonical momenta are

$$P_\Lambda = \frac{\partial \mathcal{L}}{\partial \dot{\Lambda}} = \frac{\Lambda R (N^r R' - \dot{R})}{N \sqrt{h}}, \quad (4)$$

$$P_R = \frac{\partial \mathcal{L}}{\partial \dot{R}} = \frac{R [C N^{\phi'} + \Lambda \{(\Lambda N^r)' - \dot{\Lambda}\}]}{N \sqrt{h}}, \quad (5)$$

$$P_C = \frac{\partial \mathcal{L}}{\partial \dot{C}} = -\frac{(N^r C)' + R^2 N^{\phi'} - \dot{C}}{2N \sqrt{h}}, \quad (6)$$

and the Hamiltonian and the momentum functions are defined as

$$H = \sqrt{h} \left(-\frac{P_\Lambda P_R}{\Lambda R} + P_C^2 + \frac{R'^2 + R R''}{h} - \frac{R R' h'}{2h^2} + \lambda \right), \quad (7)$$

$$H_r = R' P_R - C P_C' - \Lambda P_\Lambda', \quad (8)$$

$$H_\phi = -\left(\frac{C}{\Lambda} P_\Lambda + R^2 P_C \right)'. \quad (9)$$

In the expression of canonical momenta in Eq. (4)-(6), $\mathcal{L}$ is the Lagrangian density and $h = \Lambda^2 R^2 - C^2$ is the determinant of the spatial part of metrics.

It is essential to introduce the local conservation quantities, the angular momentum J and the mass function M as follows

$$J := -2 \int dr H_\phi = 2 \left(\frac{C}{\Lambda} P_\Lambda + R^2 P_C \right), \quad (10)$$

$$\begin{aligned} M &:= - \int dr \left(\frac{R R'}{\sqrt{h}} H + \frac{P_\Lambda}{\Lambda} H_r + P_C H_\phi \right) \\ &= \frac{1}{2} \left(P_\Lambda^2 + \frac{2C P_\Lambda P_C}{\Lambda} + R^2 P_C^2 - \frac{R^2 R'^2}{h} - \lambda R^2 + 1 \right). \end{aligned} \quad (11)$$

The angular momentum function J could be derived as a conservative quantity for a infinitesimal continuous rotation by Noether's Theorem, and the mass

function M could be derived by using the form of J and the transformation method from the canonical data which was considered by Fischler, Morgan and Polchinski [15] and by Kuchař [16]. The physical meaning of the mass function in the 3+1 dimensional space-time is studied by Nambu and Sasaki [17]. These local conserved quantities are essential in defining and solving the quantum theory in spherically symmetric gravity.

Before going to the detail of the quantum theory, we make transformation from old variables Λ, R and C into new variables by

$$\begin{pmatrix} \Lambda \\ R \\ C \end{pmatrix} \longrightarrow \begin{pmatrix} \bar{\Lambda} \\ \bar{R} \\ \bar{C} \end{pmatrix} = \begin{pmatrix} \sqrt{\Lambda^2 - C^2 R^{-2}} \\ R \\ C R^{-2} \end{pmatrix}. \quad (12)$$

The corresponding momenta are transformed as

$$\begin{pmatrix} P_\Lambda \\ P_R \\ P_C \end{pmatrix} \longrightarrow \begin{pmatrix} P_{\bar{\Lambda}} \\ P_{\bar{R}} \\ P_{\bar{C}} \end{pmatrix} = \begin{pmatrix} \sqrt{\Lambda^2 - C^2 R^{-2}} \Lambda^{-1} P_\Lambda \\ C^2 \Lambda^{-1} R^{-3} P_\Lambda + P_R + 2C R^{-1} P_C \\ C \Lambda^{-1} P_\Lambda + R^2 P_C \end{pmatrix}. \quad (13)$$

These new variables will be used extensively in the following calculation.

3 Quantum Theory

Next we proceed to the quantum theory in the Schrödinger picture. The momentum operators are expressed in the new variables in Eqs.(12) and (13) as

$$\hat{P}_{\bar{\Lambda}}(r) := -i \frac{\delta}{\delta \bar{\Lambda}(r)}, \quad \hat{P}_{\bar{R}}(r) := -i \frac{\delta}{\delta \bar{R}(r)}, \quad \hat{P}_{\bar{C}}(r) := -i \frac{\delta}{\delta \bar{C}(r)}. \quad (14)$$

The notation hat denotes the quantized operator in the followings. According to the Dirac approach, the constraints are treated as operator restriction on the wave function Ψ :

$$\hat{H}\Psi = 0, \quad \hat{H}_r\Psi = 0, \quad \hat{H}_\phi\Psi = 0. \quad (15)$$

The first equation is the Wheeler-DeWitt equation and others are the momentum constraint equations.

Our strategy is to solve the eigenvalue equation for $\hat{J}$, $\hat{M}$ and the constraint equation for $\hat{H}_r$ step by step instead of solving the constraint equations (Eq. (15)).

Step 1: Angular momentum eigenequation

The eigenvalue equation of the local angular momentum (Eqs. (10) and (13)) with the eigenvalue j

$$\hat{J}\Psi = \hat{P}_{\bar{C}}\Psi = j\Psi \quad (16)$$

is satisfied by the eigenfunction in the form

$$\Psi = e^{ij\Phi} u(\bar{\Lambda}, \bar{R}) , \quad (17)$$

with the variable

$$\Phi = \int dr \bar{C}(r) . \quad (18)$$

Step 2: Momentum constraint equation

The radial momentum constraint equation

$$\hat{H}_r\Psi = (\bar{R}'\hat{P}_{\bar{R}} - \bar{\Lambda}(\hat{P}_{\bar{\Lambda}})')e^{ij\Phi}u(\bar{\Lambda}, \bar{R}) = 0 , \quad (19)$$

restricts the functional form of the wave function to

$$\Psi = e^{ij\Phi} u(Z) , \quad (20)$$

where we introduce variable Z

$$Z = \int dr \bar{\Lambda} f(\bar{R}, \chi) = \int dr \int^{\bar{\Lambda}(r)} d\bar{\Lambda} \bar{f}(\bar{R}, \chi) , \quad (21)$$

with the definition

$$\chi := \bar{R}'^2 \bar{\Lambda}^{-2} . \quad (22)$$

The function f and $\bar{f}$ are arbitrary functions and are related each other by

$$f(\bar{R}, \chi) = - \int^\chi d\chi \frac{\bar{f}(\bar{R}, \chi)}{2\chi} . \quad (23)$$

Step 3: Mass equation

The mass operator $\hat{M}$ consists of the integration of the linear combination of constraints with respect to radial coordinate r in Eq.(11) and is imposed on the state vector as

$$(\hat{M} - m)\Psi = 0 , \quad (24)$$

where m is an integration constant. Three operators $\hat{J}$, $\hat{M}$ and $\hat{H}_r$ should form a closed algebra in order to ensure the consistency among three wave equations (16), (19) and (24):

$$[\hat{J}(r), \hat{J}(r')] = 0 , \quad (25)$$

$$[\hat{J}(r), \hat{H}_r(r')] = i\hat{J}'(r)\delta(r - r') , \quad (26)$$

$$[\hat{J}(r), \hat{M}(r')] = 0 , \quad (27)$$

$$[\hat{H}_r(r), \hat{H}_r(r')] = i(\hat{H}_r(r)\delta'(r - r') - (r \leftrightarrow r')) , \quad (28)$$

$$[\hat{H}_r(r), \hat{M}(r')] = i\hat{M}'(r)\delta(r - r') , \quad (29)$$

$$[\hat{M}(r), \hat{M}(r')] = 0 . \quad (30)$$

The factor ordering of mass operator $\hat{M}$ is determined by this requirement and is expressed as

$$\hat{M} - m = \frac{1}{2}A\hat{P}_\Lambda A^{-1}\hat{P}_\Lambda + \frac{1}{2}(-\chi + \hat{F}(\bar{R})) , \quad (31)$$

where χ is in Eq.(22) and

$$\hat{F}(\bar{R}) = 1 - 2m - \lambda \bar{R}^2 + \frac{1}{4} \hat{J}^2 \bar{R}^{-2} . \quad (32)$$

The ordering factor A is introduced to take account of the factor ordering ambiguity and is expressed by the product of two functions as

$$A = A_Z(Z) \bar{A}(\bar{R}, \chi) , \quad (33)$$

where A_Z and $\bar{A}$ are arbitrary functions with respect to the variable Z and $\bar{R}$, χ respectively. We choose one of these function $\bar{A}$ as

$$\bar{A} = \frac{\delta Z}{\delta \Lambda} = \bar{f} = \sqrt{\chi - F_j(\bar{R})} , \quad (34)$$

where

$$F_j(\bar{R}) := \hat{F} |_{\hat{j}=j} (\bar{R}) . \quad (35)$$

Then using the mass operator for each eigenvalue of angular momentum j

$$\hat{M}_j := \hat{M} |_{\hat{j}=j} , \quad (36)$$

the mass equation

$$\hat{M}_j u_{j,m}(Z) = m u_{j,m}(Z) , \quad (37)$$

can reduce to the equation with respect to Z

$$\frac{d^2 u_{j,m}(Z)}{dZ^2} - A_Z^{-1} \frac{\delta A_Z}{\delta Z} \frac{du_{j,m}(Z)}{dZ} + u_{j,m}(Z) = 0. \quad (38)$$

If we choose the remaining ordering factor as $A_Z = Z^{2\nu-1}$, the above equation becomes the Bessel equation

$$\frac{d^2 u_{j,m}(Z)}{dZ^2} - \frac{2\nu - 1}{Z} \frac{du_{j,m}(Z)}{dZ} + u_{j,m}(Z) = 0, \quad (39)$$

and the solution is

$$u_{j,m}^{(\nu)}(Z) = Z^\nu [b_1 H_\nu^{(1)}(Z) + b_2 H_\nu^{(2)}(Z)], \quad (40)$$

where $H_\nu(Z)$ is the Hankel function and b_1 and b_2 are constant coefficients. In conclusion, the quantum wave function becomes

$$\Psi(Z) = e^{ij\Phi} u_{j,m}^{(\nu)}(Z), \quad (41)$$

where the argument Z is expressed using Eqs. (21) and (34) as

$$\begin{aligned} Z &= \int dr \int^{\bar{\Lambda}(r)} d\bar{\Lambda} \sqrt{\chi - F_j(\bar{R})} \\ &= \int dr \left(\bar{\Lambda} \sqrt{\chi - F_j(\bar{R})} - \bar{R}' \ln \left| \frac{\sqrt{\chi} + \sqrt{\chi - F_j(\bar{R})}}{\sqrt{|F_j(\bar{R})|}} \right| \right), \end{aligned} \quad (42)$$

where χ and F_j are given in Eqs. (22) and (35).

The solution in Eq. (41) is shown to satisfy the original constraint equations (15) automatically, because the operators $\hat{J}$ and $\hat{M}$ are expressed by the linear combination of the original constraints (see Eqs. (10) and (11)). The closure of the algebras among original constraints H , H_r and H_ϕ is ensured in a similar way.

In the final part of this section, we comment on other choice of factor ordering for A_Z . For example, if we choose $A_Z = e^{2Z} Z^{-\sigma} (Z - 1)^\sigma$, we obtain the wave function of the hypergeometric function. The factor ordering ambiguity appears in our approach, because the factor ordering is determined by requiring a closed algebra among constraint operators and not requiring to define a inner product among state vectors.

4 De Broglie-Bohm Interpretation

We obtained the analytic wave function of the 2+1 dimensional gravity theory. The wave function is the functional of the metrics and its physical and geometrical meaning is not clear. We adopt the de Broglie-Bohm (dBB) interpretation in order to extract geometrical information from the wave function. For this purpose, we choose only the 2nd kind Hankel function in Eq. (40) as the wave function which satisfies the Vilenkin boundary condition [11]. Then the wave function in the polar coordinates is written as

$$\Psi(Z) = Z^\nu H_\nu^{(2)}(Z) = |\Psi| e^{i\Theta}(Z) . \quad (43)$$

In the dBB interpretation, the momenta are defined by the gradient of the phase of the wave function as

$$P_{\bar{\Lambda}} = \frac{\delta\Theta}{\delta\bar{\Lambda}} , \quad P_{\bar{R}} = \frac{\delta\Theta}{\delta\bar{R}} , \quad P_{\bar{C}} = \frac{\delta\Theta}{\delta\bar{C}} . \quad (44)$$

Inserting the original expressions of the momenta in Eqs. (4)- (6) and (13) into Eq. (44), we obtain the differential equations for the metrics with respect to the time and space coordinates as

$$\frac{1}{N}(N^r \bar{R}' - \dot{\bar{R}}) = \bar{f} \frac{d\Theta}{dZ} , \quad (45)$$

$$\frac{1}{N}((\bar{\Lambda} N^r)' - \dot{\bar{\Lambda}}) = \frac{\bar{\Lambda}}{\bar{R}'} \bar{f}' \frac{d\Theta}{dZ} , \quad (46)$$

$$-\frac{\bar{R}^3}{N\bar{\Lambda}}((N^r \bar{C})' + (N^\phi)' - \dot{\bar{C}}) = j . \quad (47)$$

In the dBB interpretation, the quantum effect is expressed by the factor

$$(n_{\text{dBB}})^{-1} := -\frac{d\Theta}{dZ} = \frac{2}{\pi Z |H_\nu^{(2)}(Z)|^2} . \quad (48)$$

The limit $n_{\text{dBB}} \rightarrow 1$ is the classical limit, because the dBB equations (45),(46) and (47) reduce to the Einstein equation in this limit. The metrics solving these dBB equations include quantum effects through the factor n_{dBB} .

In order to integrate the Eqs. (45)-(47), we fix a part of gauge freedom of the general coordinate transformation as

$$N^r = N^\phi = 0 . \quad (49)$$

Then the metric $\bar{C}$ is determined from Eq. (47) as

$$\dot{\bar{C}} = \frac{N\bar{\Lambda}}{\bar{R}^3} j . \quad (50)$$

We can derive the relation free from quantum effects taking the ratio of Eq. (46) over Eq. (45)

$$\frac{\dot{\bar{\Lambda}}}{\bar{\Lambda}} = \frac{\dot{\bar{R}}}{\bar{R}'} \frac{\bar{f}'}{\bar{f}} . \quad (51)$$

We show some examples of the solution by the dBB interpretation in the followings.

Example 1: BTZ black hole

If we further restrict the solution by the conditions as

$$\bar{R}' = 0 \quad \text{and} \quad \dot{\bar{R}} = 1 , \quad (52)$$

we have the relation $\bar{R} = t$. Then we find

$$ds^2 = \frac{1}{F_j} n_{\text{dBB}}^2 dt^2 - F_j dr^2 + t^2 (\bar{C} dr + d\phi)^2 , \quad (53)$$

from Eqs. (45) and (46) where $F_j = 1 - 2m + j^2/(4t^2) - \lambda t^2$. The classical limit of the metrics under the space-time transformation $t \leftrightarrow r$, $\bar{C} \leftrightarrow N^\phi$, represents

the BTZ black [2] .

Example 2: de Sitter universe

Next we show the example of de Sitter universe. We set $N = 1$ and $m = j = 0$ and we put the ansatz for metrics as

$$\bar{R} = a(t)r \quad \text{and} \quad \bar{\Lambda} = b(t)/\sqrt{1 - Kr^2} , \quad (54)$$

where $K = 0$ or 1 represent the open or closed universe respectively. From Eq. (51), we obtain $a(t) = b(t)$ and from Eq. (45) we obtain the equation for the scale factor of the universe as

$$\dot{a} = \sqrt{\lambda a^2 - K} \, n_{\text{dBB}}^{-1} . \quad (55)$$

The corresponding classical solutions (put $n_{\text{dBB}} = 1$ in Eq. (55)) are open or closed de Sitter universe

$$a = \begin{cases} \exp(\sqrt{\lambda}t) & \text{for } K = 0 \\ \lambda^{-1/2} \cosh(\sqrt{\lambda}t) & \text{for } K = 1 . \end{cases} \quad (56)$$

The classical closed de Sitter universe is created with a finite size at $t = 0$ and expands exponentially as $t \gg 0$. A special interest is the closed de Sitter universe because the finite quantum effect can be estimated.

5 Evaluation of Quantum Effect of de Sitter universe

In this section, we investigate the quantitative evaluation of quantum effects for the metrics of de Sitter universe which were studied in the previous section. We note that the quantum effect does not appear for a open de Sitter universe. This is because the argument Z becomes infinity as its spatial integration region is

infinite and therefore the factor n_{dBB} becomes one, which indicates the classical limit. The finite quantum effect is evaluated only for the case of closed de Sitter universe as was explained in the previous section:

$$\bar{R} = a(t)r \quad \text{and} \quad \bar{\Lambda} = a(t)/\sqrt{1-r^2}, \quad (57)$$

where the scale factor is determined by the dBB equation (55) with the relation (48) as

$$\dot{a} = \sqrt{\lambda a^2 - 1} \frac{2}{\pi Z |H_\nu^{(2)}(Z)|^2}. \quad (58)$$

To evaluate this equation we should first integrate with respect to r and obtain the expression of the argument Z as a function of the scale factor $a(t)$. It is expressed in the classical region ($\sqrt{\lambda}a \geq 1$) and the quantum tunneling region ($0 \leq \sqrt{\lambda}a \leq 1$) as

$$Z = \begin{cases} a\sqrt{\lambda a^2 - 1} - \frac{1}{\sqrt{\lambda}} \tanh^{-1} \frac{\sqrt{\lambda a^2 - 1}}{\sqrt{\lambda}a} & (\text{for } \sqrt{\lambda}a \geq 1), \\ i \left(a\sqrt{1 - \lambda a^2} - \frac{1}{\sqrt{\lambda}} \tan^{-1} \frac{\sqrt{1 - \lambda a^2}}{\sqrt{\lambda}a} \right) & (\text{for } 0 \leq \sqrt{\lambda}a < 1), \end{cases} \quad (59)$$

The criterion of analytic continuation from the classical region to the quantum tunneling region is that the value of Z^2 is connected smoothly at boundary point $\sqrt{\lambda}a = 1$, because Z takes the real value for the classical region and the imaginary value for the quantum region. The value of the argument Z^2 is shown in Fig. 1.

Now we turn to the evaluation of the scale factor in Eq. (58). We note that at the boundary $\sqrt{\lambda}a = 1$ the rate of change of the scale factor $\dot{a}(t)$ is zero for the classical limit ($n_{\text{dBB}} = 1$) and for $|\nu| > 1/3$ and infinity for $|\nu| < 1/3$,




Figure 1: The argument Z^2 as a function of the scale factor a given by Eq. (59) with the cosmological constant $\lambda = 1/2$

where ν is the index of the Hankel function. The interesting tunneling effect is obtained for the case of index of the Hankel function $\nu = 1/3$, where the rate of change of the scale factor takes a finite value at the boundary:

$$\lim_{\sqrt{\lambda}a \rightarrow 1} \dot{a} = \lim_{\sqrt{\lambda}a \rightarrow 1} \sqrt{\lambda a^2 - 1} \frac{2}{\pi Z |H_{1/3}^{(2)}(Z)|^2} = \frac{\lambda^{1/6} 3^{4/3} \Gamma(2/3)^2}{4\pi}. \quad (60)$$

The case for $\nu = -1/3$ gives the same result because of the identity: $H_{-\nu}^{(2)}(Z) = \exp(-i\pi\nu)H_{\nu}^{(2)}(Z)$. Then the scale factor both in the classical region and the quantum region is determined by

$$\dot{a}(t) = \begin{cases} \frac{2\sqrt{\lambda a^2 - 1}}{\pi Z |H_{1/3}^{(2)}(Z)|^2} & (\text{for } \sqrt{\lambda}a \geq 1), \\ \frac{2i\sqrt{1 - \lambda a^2}}{\pi Z |H_{1/3}^{(2)}(Z)|^2} & (\text{for } 0 \leq \sqrt{\lambda}a < 1), \end{cases} \quad (61)$$

where the argument Z is given in Eq. (59). From this equation the rate of change of the scale factor $\dot{a}(t)$ as a function $a(t)$ is obtained and shown in Fig. 2 with $\lambda = \frac{1}{2}$. The scale factor as a function of t is obtained by integrating




Figure 2: The rate of change of the scale factor $\dot{a}(t)$ as a function $a(t)$. (A) The classical case: $\dot{a} = \sqrt{\lambda a^2 - 1}$ and (B) The quantum case: $\dot{a}$ in Eq. (58) with the cosmological constant $\lambda = 1/2$.

Eq.(61) numerically and the result is shown in Fig. 3.

Here we can see from Figs. 2 and 3 that the quantum scale factor (denoted by a letter B) approaches to the classical scale factor (denoted by a letter A) asymptotically ($a(t) \rightarrow \infty$), which was required by the Vilenkin's boundary condition. The quantum scale factor extend in the classically forbidden region ($t \leq 0$) as a real time quantum tunneling effect. This is one of characteristic features of the de Broglie-Bohm approach and are compared with other approaches: the WKB approach [11] or the path integral approach [12]. A similar analysis was done by Horiguchi [18] in 3+1 dimension, in which metrics were treated as function of only time t while we have treated more generally spherically symmetric metrics as function of both time and radial coordinates.




Figure 3: The scale factor $a(t)$ for the closed de Sitter universe. (A) The classical scale factor $a(t) = \lambda^{-1/2} \cosh(\sqrt{\lambda}t)$, (B) The numerical solution of Eq. (61), which includes quantum effects. The cosmological constant is set to $\lambda = 1/2$ and the time is normalized as $a = 1/\sqrt{\lambda}$ at $t = 0$.

6 Summary

We have studied the 2+1 dimensional spherically symmetric gravity theory and obtained the following results.

- (1) Quantum theory is defined through local conserved quantities $\hat{J}(r)$ and $\hat{M}(r)$.
- (2) The de Broglie-Bohm interpretation is applied for the analytic wave function of universe.
- (3) The differential equations of the dBB interpretation are solved and black hole metrics as well as expanding universe metrics are obtained after fixing the coordinate conditions.
- (4) Especially we have evaluated the quantum effects of the closed de Sitter universe as the real time tunneling in the classically forbidden region occurs through the de Broglie-Bohm interpretation. It is interesting to compare with other approaches that the universe creates form nothing through imaginary time tunneling using WKB approximation by Vilenkin [11] and the path integral

method by Hartle and Hawking [12].

There are some other interesting topics characteristic in 2+1 dimensional gravity. Localized sources can affect global geometrical structure which is expressed as topological conical singularity [1]. Other interesting topological solutions are torus solutions considered by Hosoya and Nakano [19]. It may be interesting to make the de Broglie-Bohm interpretation for the conical singularity in the closed universe and torus solutions in 2+1 dimensional gravity because the evaluation of the quantum effects are expected to be finite and the relation between the quantum solutions and the classical solutions becomes clear. These are our future problems.

7 Acknowledgements

We would like to thank T.Takahashi for useful discussions and numerical calculations.

References

- [1] S. Deser, R. Jackiw and G.t' Hooft, Ann. Phys. **152**, 220(1984)
- [2] M. Banados, C. Teitelboim and J. Zanelli , Phys. Lett. **69**, 1849 (1992).
- [3] Y. Yoshida and T. Kubota, Phys. Rev. **D60**, 044013 (1999).
- [4] E. Witten, Nucl. Phys. **B311**, 46(1988).
- [5] S.W. Hawking, Commun. Math. Phys. **43**, 199(1975).
- [6] A. Achúcarro and P.K. Townsend, Phys. Lett. **B180**, 89(1986).

- [7] S. Deser and X. Xiang, Phys. Lett. **B263** 39(1991).
- [8] D. Bohm, Phys. Rev. **85**, 166(1952); **85**, 180(1952).
- [9] J.S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, 1987).
- [10] P.R. Holland, *Quantum Theory of Motion* (Cambridge University Press, Cambridge, 1993). (Cambridge University Press, Cambridge, 1987).
- [11] A. Vilenkin, Phys. Lett. **117B**, 25(1982); Phys. Rev. **D27**, 2848(1983); Phys. Rev. **D37**, 888(1988).
- [12] J.B. Hartle and S. Hawking, Phys. Rev. **D28**, 2960(1983).
- [13] M. Kenmoku, H. Kubotani, E. Takasugi and Y. Yamazaki , Phys. Rev. **D59**, 124004 (1999); Prog. Theor. Phys. **105**, 897 (2001).
- [14] R. Arnowitt, S. Deser and C.W. Misner, in Gravitation: An Introduction to Current Research, edited by L.Witten (Wiley, New York, 1962).
- [15] W. Fishler, D. Morgan and J. Polchinski, Phys. Rev. **D42**, 4042(1990).
- [16] K.V. Kuchař, Phys. Rev. **D50**, 3961 (1994).
- [17] Y. Nambu and M. Sasaki, Prog. Theor. Phys. **79**, 96 (1988).
- [18] T. Horiguchi, Mod. phys. Lett. **A9**, 1429 (1994).
- [19] A. Hosoya and K. Nakano, Prog. Theor. Phys. **84**, 739 (1990); Class. Quantum Grav. **7**, 163 (1990).